

New Study Points to Proteins and Salt for Life's Origin


A structural biologist at the Florida State University College of Medicine has made discoveries that could lead scientists a step closer to understanding how life [first emerged](#) on Earth billions of years ago.

Professor Michael Blaber and his team produced data supporting the idea that 10 [amino acids](#) believed to exist on Earth around 4 billion years ago were capable of forming foldable proteins in a high-salt (halophile) environment. Such proteins would have been capable of providing metabolic activity for the first living organisms to emerge on the planet between 3.5 and 3.9 billion years ago.

The results of Blaber's three-year study, which was built around investigative techniques that took more than 17 years to develop, are published in the journal *Proceedings of the National Academy of Sciences*.

The first living [organisms](#) would have been microscopic, cell-like organizations capable of replicating and adapting to environmental conditions — a humble beginning to life on Earth.

"The current paradigm on the emergence of life is that RNA came first and in a high-temperature environment," Blaber said. "The data we are generating are much more in favor of a protein-first view in a halophile environment."



The two enantiomers of the amino acid alanine (D-Alanine and L-Alanine). Amino acids are essential for life as we know it, and are the building blocks of proteins in living cells. Credit: [Wikimedia Commons](#)

Without the ability to fold, proteins would not be able to form the precise structures essential for functions that sustain life as we know it. Folding allows proteins to take on a globular shape through which they can interact with other proteins, perform specific chemical reactions, and adapt to enable organisms to exploit a given environment.

"There are numerous niches that life can evolve into," Blaber said. "For example, extremophiles are organisms that exist in high temperatures, high [acidity](#), extreme cold, extreme pressure and extreme salt and so on. For life to exist in such environments it is essential that proteins are able to adapt in those conditions. In other words, they have to be able to fold."



Professor Michael Blaber of Florida State University. Credit: [Florida State University](#)

The widely accepted view among scientists is that RNA, found in all living cells, would have likely represented the first molecules of life, hypothesizing an "[RNA-first](#)" view of the origin of living systems from non-living molecules. Blaber's results indicate that the set of amino acids produced by simple chemical processes contains the requisite information to produce complex folded proteins, which supports an opposing "protein-first" view.

Another prevailing view holds that a high-temperature (thermophile) environment, such as deep-ocean thermal vents, may have been the breeding ground for the origin of life.

"The halophile, or salt-loving, environment has typically been considered one that life adapted into, not started in," Blaber said. "Our study of the [prebiotic amino acids](#) and protein design and folding suggests the opposite."

Comet and meteorite fragments, like those that recently struck in the Urals region of Russia, have provided evidence regarding the arrival of amino acids on Earth. Such fragments predate the earth and would have been responsible for delivering a set of 10 prebiotic (before life) amino acids, whose origins are in the formation of our solar system.

Today the human body uses 20 common amino acids to make all its [proteins](#). Ten of those emerged through biosynthetic pathways — the way living systems evolve. Ten — the prebiotic set — can be made by chemical reactions without requiring any living system or biosynthetic pathway.


Scientific evidence exists to support many elements in theories of abiogenesis (the emergence of life), including the time frame (around 3.5 to 3.9 billion years ago) and the conditions on Earth and in its atmosphere at that time. Earth would have been made up of volcanic land masses (the beginning of the formation of continents), salty oceans and freshwater ponds, along with a hot (around 80 degrees Celsius) and steamy atmosphere comprising carbon dioxide and nitrogen. Oxygen would have come later as a by-product of green plant life and bacteria that emerged.

Using a technique called top-down symmetric deconstruction, Blaber's lab has been able to identify small peptide building blocks capable of spontaneous assembly into specific and complex protein architectures. His recent work explored whether such building blocks can be comprised of only the 10 prebiotic amino acids and still fold.

His team has achieved foldability in [proteins](#) down to 12 amino acids — about 80 percent of the way to proving his hypothesis.

If Blaber's theory holds, scientists may refocus where they look for evidence in the quest to understand where, and how, life began.

"Rather than a curious niche that life evolved into, the halophile environment now may take center stage as the likely location for key aspects of abiogenesis," he said. "Likewise, the role of the formation of proteins takes on additional importance in the earliest steps in the beginnings of life on Earth."



The single-helix structure of RNA. Many scientists believe that the origin of life was dependent on RNA forming in a high-temperature environment. However, the new study challenges the RNA-first view. Credit: Darryl Leja/Access Excellence